

Set-Theoretical Kripke-Style Semantics for an Extension of HpsUL, CnHpsUL*

Eunsuk Yang

【Abstract】 This paper deals with non-algebraic Kripke-style semantics, i.e, set-theoretical Kripke-style semantics, for weakening-free *non-commutative* fuzzy logics. We first recall an extension of the pseudo-uniform based fuzzy logic **HpsUL**, **CnHpsUL***. We next introduce set-theoretical Kripke-style semantics for it.

【Key Words】 (Set-theoretical) Kripke-style semantics, Algebraic semantics, Fuzzy logic, **HpsUL**, **CnHpsUL***.

Received: Nov. 21, 2017. Revised: Jan. 29, 2018. Accepted: Jan. 5, 2018.

* This research was supported by “Research Base Construction Fund Support Program” funded by Chonbuk National University in 2017. I must thank the referees for their helpful comments.

1. Introduction

The aim of this paper is to introduce set-theoretic Kripke-style semantics for weakening-free non-commutative substructural fuzzy logic. For this, note that Yang recently introduced two kinds of (binary) Kripke-style semantics, i.e., algebraic and non-algebraic Kripke-style semantics, for logics with pseudo-Boolean (briefly, pB) and de Morgan (briefly, dM) negations in Yang (2015b). He (2014b, 2015a) further considered such semantics for logics with weak-Boolean (briefly, wB) negations, which can be regarded as paraconsistent logics. Recently, he (2016) introduced algebraic Kripke-style semantics for a weakening-free non-commutative substructural fuzzy logic, **CnHpsUL***. But he did not consider set-theoretical semantics for it. Thus, it is not clear whether this semantics works for weakening-free non-commutative substructural fuzzy logic systems.

This is a tough question because Kripke-style semantics for well-known core fuzzy systems are algebraic, but not set-theoretical. Recall some historical facts associated with this. As Yang mentioned in Yang (2014a), after introducing algebraic semantics for t-norm¹⁾ (based) logics, their corresponding algebraic Kripke-style semantics have been introduced: after Esteva and Godo introducing algebraic semantics for monoidal t-norm (based) logics in Esteva & Godo (2001), their corresponding algebraic Kripke-style semantics were introduced in Montagna & Ono

¹⁾ T-norms are commutative, associative, increasing, binary functions with identity 1 on the real unit interval [0,1].

(2002), Montagna & Sacchetti (2003; 2004), and Diaconescu & Georgescu (2007). Furthermore, algebraic semantics and corresponding algebraic Kripke-style semantics for core fuzzy logic systems based on more general structures have been introduced: after Hájek introducing algebraic semantics for *non-commutative* pseudo-t-norm (based) logics in Hájek (2003a; 2003b), one corresponding algebraic Kripke-style semantics for the pseudo-t-norm (based) logic **psMTL**[†] was introduced in Diaconescu (2010). After Metcalfe and Montagna introducing algebraic semantics for *weakening-free* uninorm (based) logics in Metcalfe & Montagna (2007), their corresponding algebraic Kripke-style semantics were introduced in Yang (2012; 2014a). After Wang (2013) introducing algebraic semantics for **CnHpsUL***, the **HpsUL*** with n-potency, its corresponding algebraic Kripke-style semantics was introduced in Yang (2016).

Then, these facts raise the following interesting question:

- Can we introduce set-theoretical Kripke-style semantics for core fuzzy systems, in particular **CnHpsUL***?

The answer to the question is positive in the sense that we can provide such Kripke-style semantics for **CnHpsUL***. For this, first, in Section 2 we recall the system **CnHpsUL***. In Section 3, we introduce the other kind of binary relational Kripke-style semantics, non-algebraic set-theoretical Kripke-style semantics, for **CnHpsUL***.

For convenience, we shall adopt the notation and terminology

similar to those in Cintula (2006), Metcalfe & Montagna (2007), Montagna & Sacchetti (2003; 2004), and Yang (2012; 2014a; 2016), and we assume reader familiarity with them (along with results found therein).

2. Preliminaries: The logic **CnHpsUL***

We base **CnHpsUL*** on a countable propositional language with formulas Fm built inductively as usual from a set of propositional variables VAR , binary connectives \rightarrow , \ddagger , $\&$, \wedge , \vee , and constants \mathbf{T} , \mathbf{F} , \mathbf{t} ²⁾, with a defined connective:

$$\text{df1. } \phi \leftrightarrow \psi := (\phi \rightarrow \psi) \wedge (\psi \rightarrow \phi).$$

We moreover define ϕ^n_t as $\phi_t \& \cdots \& \phi_t$, n factors, where $\phi_t := \phi \wedge \mathbf{t}$. For the remainder we shall follow the customary notation and terminology. We use the axiom systems to provide a consequence relation.

Definition 2.1 (i) (Metcalfe et al. (2009), Tsinakis & Blount (2003), Wang (& Zhao) (2009; 2013)) **HpsUL** consists of the following axiom schemes and rules:

- A1. $\phi \rightarrow \phi$ (self-implication, SI)
- A2. $(\phi \wedge \psi) \rightarrow \phi$, $(\phi \wedge \psi) \rightarrow \psi$ (\wedge -elimination, \wedge -E)
- A3. $((\phi \rightarrow \psi) \wedge (\phi \rightarrow \chi)) \rightarrow (\phi \rightarrow (\psi \wedge \chi))$ (\wedge -introduction, \wedge -I)
- A4. $\phi \rightarrow (\phi \vee \psi)$, $\psi \rightarrow (\phi \vee \psi)$ (\vee -introduction, \vee -I)

²⁾ The constant \mathbf{t} corresponds to the least designated element.

A5. $((\phi \rightarrow \chi) \wedge (\psi \rightarrow \chi)) \rightarrow ((\phi \vee \psi) \rightarrow \chi)$ (\vee -elimination, \vee -E)

A6. $\phi \rightarrow \mathbf{T}$ (verum ex quolibet, VE)

A7. $\mathbf{F} \rightarrow \phi$ (ex falso quadlibet, EF)

A8. \mathbf{t}

A9. $\phi \rightarrow (\mathbf{t} \rightarrow \phi)$

A10. $(\psi \rightarrow \chi) \rightarrow ((\phi \rightarrow \psi) \rightarrow (\phi \rightarrow \chi))$ (prefixing, PF)

A11. $\phi \rightarrow ((\phi \Downarrow \psi) \rightarrow \psi)$

A12. $(\phi \Downarrow (\psi \rightarrow \chi)) \rightarrow (\psi \rightarrow (\phi \Downarrow \chi))$

A13. $\psi \rightarrow (\phi \rightarrow (\phi \& \psi))$

A14. $(\psi \rightarrow (\phi \rightarrow \chi)) \rightarrow ((\phi \& \psi) \rightarrow \chi)$

A15. $((\psi \Downarrow \psi) \& (\psi \rightarrow \phi)) \rightarrow (\psi \Downarrow \phi)$

A16. $(\phi_{\mathbf{t}} \& \psi_{\mathbf{t}}) \rightarrow (\phi \wedge \psi)$

A17. $(\phi \vee \psi)_{\mathbf{t}} \rightarrow (\phi_{\mathbf{t}} \vee \psi_{\mathbf{t}})$ (prelinearity, PRL1)

A18. $(\chi \rightarrow (((\phi \vee \psi) \rightarrow \phi) \& \chi)) \vee (\chi \Downarrow (\chi \& ((\phi \vee \psi) \rightarrow \psi)))$ (PRL2)

$\phi \rightarrow \psi, \phi \vdash \psi$ (mp)

$\phi \vdash \phi_{\mathbf{t}}$ (adj_t)

$\phi \vdash \psi \rightarrow (\phi \& \psi)$ (pn \rightarrow)

$\phi \vdash \psi \Downarrow (\psi \& \phi)$ (pn \Downarrow).

(ii) (Wang (2013)) **CnHpsUL** is **HpsUL** plus $\phi^n \leftrightarrow \phi^{n-1}$, for $2 \leq n$ (n-potency, nP).

(iii) (Wang (2013)) **CnHpsUL*** is **CnHpsUL** plus $(\phi \& \psi) \rightarrow \mathbf{t} \vdash (\psi \& \phi) \rightarrow \mathbf{t}$ ³⁾ (weak commutativity, WCM).

Proposition 2.2 (Cintula, Horčík, & Noguera (2013), Yang (2016)) **CnHpsUL*** proves:

(1) $\phi \rightarrow \psi \vdash \phi \Downarrow \psi, \phi \Downarrow \psi \vdash \phi \rightarrow \psi$

³⁾ Note that we may instead take $(\phi \Downarrow \mathbf{t}) \rightarrow (\phi \Downarrow \mathbf{t})$

- (2) $\phi \rightarrow (\psi \rightarrow \chi) \vdash (\psi \& \phi) \rightarrow \chi$ (residuation1, Res1)
- (3) $\phi \rightarrow (\psi \Downarrow \chi) \vdash (\phi \& \psi) \rightarrow \chi$ (Res1 \Downarrow)
- (4) $(\psi \& \phi) \rightarrow \chi \vdash \phi \rightarrow (\psi \rightarrow \chi)$ (Res2)
- (5) $(\phi \& \psi) \rightarrow \chi \vdash \phi \rightarrow (\psi \Downarrow \chi)$ (Res2 \Downarrow)
- (6) $\phi \rightarrow \psi \vdash (\chi \& \phi) \rightarrow (\chi \& \psi), \phi \rightarrow \psi \vdash (\phi \& \chi) \rightarrow (\psi \& \chi)$
- (7) $(\mathbf{t} \& \phi) \leftrightarrow \phi \leftrightarrow (\phi \& \mathbf{t})$
- (8) $(\phi \& (\phi \rightarrow \psi)) \rightarrow \psi$.

A *theory* over $\mathbf{CnHpsUL}^*$ is a set T of formulas. A *proof* in a sequence of formulas whose each member is either an axiom of $\mathbf{CnHpsUL}^*$ or a member of T or follows from some preceding members of the sequence using the two rules in Definition 2.1. $T \vdash \phi$, more exactly $T \vdash_{\mathbf{CnHpsUL}^*} \phi$, means that ϕ is *provable* in T with respect to (w.r.t.) $\mathbf{CnHpsUL}^*$, i.e., there is a $\mathbf{CnHpsUL}^*$ -proof of ϕ in T .

The deduction theorem for $\mathbf{CnHpsUL}^*$ is following:

Definition 2.3 (Cintula & Noguera (2011)) Let T be a theory over $\mathbf{CnHpsUL}^*$, and ϕ, ψ formulas. L is almost (MP)-based with the set of basic deduction terms $\{\lambda_a(\star), \rho_a(\star) : a \in \text{Fm}\}$. Therefore, the following holds:

$T, \phi \vdash_L \psi$ if and only if (iff) $T \vdash \chi(\phi) \rightarrow \psi$ for some conjunction χ of iterated conjugates⁴).

A theory T is *inconsistent* if $T \vdash \mathbf{F}$; otherwise it is *consistent*.

⁴) For the notion of conjugate, see Cintula & Noguera (2011) and Yang (2016).

3. Kripke-style semantics for CnHpsUL*

We consider here set-theoretical Kripke-style semantics for CnHpsUL*.

Definition 3.1 (Yang (2016))

(i) (Operational Kripke frame) An *operational Kripke frame* is a structure $\mathbf{X} = (X, \top, \perp, t, f, \leq, *)$ such that $(X, \top, \perp, t, f, \leq, *)$ is a linearly ordered pointed bounded monoid. The elements of \mathbf{X} are called *nodes*.

(ii) (Residuated operational Kripke frame) An operational Kripke frame is said to be *residuated* if it has suprema w.r.t. $*$, i.e., for every $x, y \in X$, the sets $\{z: x * z \leq y\}$ and $\{z: z * x \leq y\}$ have suprema.

(iii) (CnHpsUL* frame) A *CnHpsUL* frame* is a residuated operational Kripke frame, where $*$ is conjunctive (i.e., $\perp * \top = \perp$) and left-continuous (i.e., whenever $\sup\{x_i : i \in I\}$ exists, $x * \sup\{x_i : i \in I\} = \sup\{x * x_i : i \in I\}$ and $\sup\{x_i : i \in I\} * x = \sup\{x_i * x : i \in I\}$).

Definition 3.2 ensures that a CnHpsUL* frame has suprema w.r.t. $*$, i.e., for every $x, y \in X$, the sets $\{z: x * z \leq y\}$ and $\{z: z * x \leq y\}$ have the suprema. \mathbf{X} is said to be *complete* if \leq is a complete order.

An *evaluation* or *forcing* on an algebraic Kripke frame is a relation \Vdash between nodes and propositional variables, and arbitrary formulas subject to the conditions below: for every

propositional variable p ,

(AHC) if $x \Vdash p$ and $y \leq x$, then $y \Vdash p$;

(min) $\perp \Vdash p$; and

for arbitrary formulas,

(t) $x \Vdash \mathbf{t}$ iff $x \leq \mathbf{t}$;

(f) $x \Vdash \mathbf{f}$ iff $x \leq \mathbf{f}$;

(\perp) $x \Vdash \mathbf{F}$ iff $x = \perp$;

(\wedge) $x \Vdash \phi \wedge \psi$ iff $x \Vdash \phi$ and $x \Vdash \psi$;

(\vee) $x \Vdash \phi \vee \psi$ iff $x \Vdash \phi$ or $x \Vdash \psi$;

($\&$) $x \Vdash \phi \& \psi$ iff there are $y, z \in X$ such that $y \Vdash \phi$, $z \Vdash \psi$, and $x \leq y * z$;

(\rightarrow) $x \Vdash \phi \rightarrow \psi$ iff for all $y \in X$, if $y \Vdash \phi$, then $y * x \Vdash \psi$;

(\Downarrow) $x \Vdash \phi \Downarrow \psi$ iff for all $y \in X$, if $y \Vdash \phi$, then $x * y \Vdash \psi$.

An evaluation or forcing on a CnHpsUL^* frame is an evaluation or forcing further satisfying that (max) for every atomic sentence p , $\{x : x \Vdash p\}$ has a maximum.

Definition 3.2 (Yang (2016))

(i) (Residuated operational Kripke model) A residuated operational is a pair (\mathbf{X}, \Vdash) , where \mathbf{X} is a residuated operational Kripke frame and \Vdash is a forcing on \mathbf{X} .

(ii) (CnHpsUL* model) A CnHpsUL* model is a pair (\mathbf{X}, \Vdash) , where \mathbf{X} is a CnHpsUL* frame and \Vdash is a forcing on \mathbf{X} . A CnHpsUL* model (\mathbf{X}, \Vdash) is said to be *complete* if \mathbf{X} is a complete frame and \Vdash is a forcing on \mathbf{X} .

Definition 3.3 (Cf. Montagna & Sacchetti (2004)) Given a residuated operational Kripke model (\mathbf{X}, \Vdash) , a node x of \mathbf{X} and a formula ϕ , we say that x *forces* ϕ to express $x \Vdash \phi$. We say that ϕ is *true* in (\mathbf{X}, \Vdash) if $t \Vdash \phi$, and that ϕ is *valid* in the frame \mathbf{X} (expressed by \mathbf{X} models ϕ) if ϕ is true in (\mathbf{X}, \Vdash) for every forcing \Vdash on \mathbf{X} .

Definition 3.4 A residuated operational Kripke frame \mathbf{X} is a CnHpsUL* frame iff all axioms of CnHpsUL* are valid in \mathbf{X} . We say that a CnHpsUL* model (\mathbf{X}, \Vdash) is a *CnHpsUL* model* if \mathbf{X} is a CnHpsUL* frame.

For soundness and completeness for CnHpsUL*, let $\vdash_{\text{CnHpsUL}^*}$ ϕ be the theoremhood of ϕ in CnHpsUL*.

Proposition 3.5 (Soundness, Yang (2016)) If $\vdash_{\text{CnHpsUL}^*}$ ϕ , then ϕ is valid in every CnHpsUL* frame.

Now we provide completeness results for CnHpsUL* using set-theoretical Kripke-style semantics. A theory T is said to be *linear* if, for each pair ϕ, ψ of formulas, we have $T \vdash \phi \rightarrow \psi$ or $T \vdash \psi \rightarrow \phi$. By a CnHpsUL*-theory, we mean a theory T

closed under rules of **CnHpsUL***. As in relevance logic, by a regular **CnHpsUL***-theory, we mean a **CnHpsUL***-theory containing all of the theorems of **CnHpsUL***. Since we have no use of irregular theories, henceforth, by a **CnHpsUL***-theory, we henceforth we mean a **CnHpsUL***-theory containing all of the theorems of **CnHpsUL***.

Moreover, where T is a linear **CnHpsUL***-theory, we define the *canonical CnHpsUL* frame* determined by T to be a structure $\mathbf{X} = (X_{\text{can}}, \top_{\text{can}}, \perp_{\text{can}}, t_{\text{can}}, f_{\text{can}}, \leq_{\text{can}}, *_{\text{can}})$, where $\top_{\text{can}} = \{\phi : T \vdash_{\text{CnHpsUL}^*} \mathbf{T} \rightarrow \phi\}$, $\perp_{\text{can}} = \{\phi : T \vdash_{\text{CnHpsUL}^*} \mathbf{F} \rightarrow \phi\}$, $t_{\text{can}} = T$, $f_{\text{can}} = \{\phi : T \vdash_{\text{CnHpsUL}^*} \mathbf{f} \rightarrow \phi\}$, X_{can} is the set of linear **CnHpsUL***-theories extending t_{can} , \leq_{can} is \supseteq restricted to X_{can} , i.e., $x \leq_{\text{can}} y$ iff $\{\phi : x \vdash_{\text{CnHpsUL}^*} \phi\} \supseteq \{\phi : y \vdash_{\text{CnHpsUL}^*} \phi\}$, and $*_{\text{can}}$ is defined as $x *_{\text{can}} y := \{\phi \ \& \ \psi : \text{for some } \phi \in x, \psi \in y\}$ satisfying groupoid properties corresponding to **CnHpsUL*** frames on $(X_{\text{can}}, t_{\text{can}}, \leq_{\text{can}})$. Note that the base t_{can} is constructed as the linear **CnHpsUL***-theory that excludes nontheorems of **CnHpsUL***, i.e., excludes ϕ such that $\not\vdash_{\text{CnHpsUL}^*} \phi$. The partial orderedness and the linear orderedness of the canonical **CnHpsUL*** frame depend on \leq_{can} restricted on X_{can} . Then, first, the following is obvious.

Proposition 3.6 A canonical **CnHpsUL*** frame is linearly ordered.

Proof: It is easy to show that a canonical L frame is partially ordered. We show that this frame is connected and so linearly

ordered. Suppose toward contradiction that neither $x \leq_{\text{can}} y$ nor $y \leq_{\text{can}} x$. Then, there are ϕ, ψ such that $\phi \in y, \phi \notin x, \psi \in x,$ and $\psi \notin y$. Note that, since t_{can} is a linear theory, $\phi \rightarrow \psi \in t_{\text{can}}$ or $\psi \rightarrow \phi \in t_{\text{can}}$. Let $\phi \rightarrow \psi \in t_{\text{can}}$ and thus $\phi \rightarrow \psi \in y$. Then, by (mp), we have $\psi \in y$, a contradiction. The case, where $\psi \rightarrow \phi \in t_{\text{can}}$, is analogous. \square

Next, we define a canonical evaluation as follows:

$$(a) \quad x \vdash_{\text{can}} \phi \text{ iff } \phi \in x.$$

This definition allows us to state the following lemmas.

Lemma 3.7 $t_{\text{can}} \vdash_{\text{can}} \phi \rightarrow \psi$ iff for all $x \in X_{\text{can}}$, if $x \vdash_{\text{can}} \phi$, then $x \vdash_{\text{can}} \psi$.

Proof: By (a), we need to show that $\phi \rightarrow \psi \in t_{\text{can}}$ iff for all $x \in X_{\text{can}}$, if $\phi \in x$, then $\psi \in x$. For the left-to-right direction, we assume $\phi \rightarrow \psi \in t_{\text{can}}$ and $\phi \in x$, and show $\psi \in x$. The definition of $*_{\text{can}}$ ensures $\phi \ \& \ (\phi \rightarrow \psi) \in x *_{\text{can}} t_{\text{can}} = x$. By Proposition 2.2 (8), we have $(\phi \ \& \ (\phi \rightarrow \psi)) \rightarrow \psi \in t_{\text{can}}$ and thus $(\phi \ \& \ (\phi \rightarrow \psi)) \rightarrow \psi \in x$. Therefore, we obtain $\psi \in x$ by (mp). We prove the other direction contrapositively. Suppose $\phi \rightarrow \psi \notin t_{\text{can}}$. We set $x_0 = \{Z : \text{there exists } X \in t_{\text{can}} \text{ and } t_{\text{can}} \vdash (\phi \ \& \ X) \rightarrow Z\}$. Clearly, $x_0 \supseteq t_{\text{can}}, \phi \in x_0$, but also $\psi \notin x_0$. (Otherwise, $t_{\text{can}} \vdash (\phi \ \& \ X) \rightarrow \psi$ and thus $t_{\text{can}} \vdash X \rightarrow (\phi \rightarrow \psi)$; therefore, since $t_{\text{can}} \vdash X$, by (mp), we have $t_{\text{can}} \vdash$

$\phi \rightarrow \psi$, a contradiction.)

Then, by the Linear Extension Property of Theorem 12.9 in Cintula, Horčík, & Noguera (2015), we have a linear theory $x \supseteq x_0$ with $\psi \notin x$; therefore $\phi \in x$ but $\psi \notin x$. \square

Lemma 3.8 (Canonical Evaluation Lemma) \Vdash_{can} is an evaluation.

Proof: We first consider the conditions for propositional variables.

For (AHC), we must show that: for every propositional variable p ,

$$\text{if } x \Vdash_{\text{can}} p \text{ and } y \leq_{\text{can}} x, \text{ then } y \Vdash_{\text{can}} p.$$

Let $x \Vdash_{\text{can}} p$ and $y \leq_{\text{can}} x$. By (a), we have $p \in x$ and $x \subseteq y$, and thus $p \in y$. Hence, by (a), we have $y \Vdash_{\text{can}} p$.

For (min), we must show that: for every propositional variable p ,

$$\perp_{\text{can}} \Vdash_{\text{can}} p.$$

By (a), we need to show that $p \in \perp_{\text{can}}$. Since $\perp_{\text{can}} = \{\phi : T \vdash_{\text{cHpsUL}^*} \mathbf{F} \rightarrow \phi\}$, $p \in \perp_{\text{can}}$.

We next consider the conditions for propositional constants \mathbf{t} , \mathbf{f} , and \mathbf{F} .

For (t), we must show that:

$$x \Vdash_{\text{can}} \mathbf{t} \text{ iff } x \leq_{\text{can}} t_{\text{can}}.$$

By (a), we need to show that $\mathbf{t} \in x$ iff $x \supseteq t_{\text{can}}$. This is obvious since $t_{\text{can}} = T$ and x is a theory extending T .

For (f), we must show that:

$$x \Vdash_{\text{can}} \mathbf{f} \text{ iff } x \leq_{\text{can}} f_{\text{can}}.$$

By (a), we need to show that $\mathbf{f} \in x$ iff $x \supseteq f_{\text{can}}$. This is obvious since $f_{\text{can}} = \{\phi : T \vdash_{\text{CnHpsUL}^*} \mathbf{f} \rightarrow \phi\}$ and x is a theory extending T .

For (\perp), we must show that:

$$x \Vdash_{\text{can}} \mathbf{F} \text{ iff } x =_{\text{can}} \perp_{\text{can}}.$$

By (a), we need to show that $\mathbf{F} \in x$ iff $x =_{\text{can}} \perp_{\text{can}}$. This is obvious since $\perp_{\text{can}} = \{\phi : T \vdash_{\text{CnHpsUL}^*} \mathbf{F} \rightarrow \phi\}$.

Now we consider the conditions for arbitrary formulas.

For (\wedge), we must show

$$x \Vdash_{\text{can}} \phi \wedge \psi \text{ iff } x \Vdash_{\text{can}} \phi \text{ and } x \Vdash_{\text{can}} \psi.$$

By (a), we need to show that $\phi \wedge \psi \in x$ iff $\phi \in x$ and $\psi \in x$. The left-to-right direction follows from (\wedge -E) and (mp). The right-to-left direction follows from (adj).

For (\vee), we must show

$$x \Vdash_{\text{can}} \phi \vee \psi \text{ iff } x \Vdash_{\text{can}} \phi \text{ or } x \Vdash_{\text{can}} \psi.$$

By (a), we need to show that $\phi \vee \psi \in x$ iff $\phi \in x$ or $\psi \in x$. The left-to-right direction follows from the fact that linear theories are also prime theories in **CnHpsUL*** (see Cintula & Noguera (2011)). The right-to-left direction follows from (\vee -I) and (mp).

For ($\&$), we must show

$$x \Vdash_{\text{can}} \phi \ \& \ \psi \text{ iff there are } y, z \in X \text{ such that } y \Vdash_{\text{can}} \phi, z \Vdash_{\text{can}} \psi, \text{ and } x = y *_{\text{can}} z.$$

By (a), we need to show that $\phi \ \& \ \psi \in x$ iff there are $y, z \in X$ such that $\phi \in y$, $\psi \in z$, and $x = y *_{\text{can}} z$. This directly follows from the definition of $*_{\text{can}}$.

For (\rightarrow), we must show

$$x \Vdash_{\text{can}} \phi \rightarrow \psi \text{ iff for all } y \in X, \text{ if } y \Vdash_{\text{can}} \phi, \text{ then } y *_{\text{can}} x \Vdash_{\text{can}} \psi.$$

By (a), we need to show that $\phi \rightarrow \psi \in x$ iff for all $y \in X$, if $\phi \in y$, then $\psi \in y *_{\text{can}} x$. For the left-to-right direction, we assume $\phi \rightarrow \psi \in x$ and $\phi \in y$, and show $\psi \in y *_{\text{can}} x$. The definition of $*_{\text{can}}$ ensures $\phi \ \& \ (\phi \rightarrow \psi) \in y *_{\text{can}} x$. Then, by Proposition 2.2 (8) and Lemma 3.7, we obtain $\psi \in y *_{\text{can}} x$. We prove the right-to-left direction contrapositively. Suppose $\phi \rightarrow \psi \notin x$. We need to construct a linear theory y such that $\phi \in y$

and $\psi \notin y \text{ *}_{\text{can}} x$. Let y_0 be the smallest regular **CnHpsUL***-theory extending t_{can} with $\{\phi\}$ and satisfying $y_0 \text{ *}_{\text{can}} x = \{Z : \text{there is } X \in x \text{ and } t_{\text{can}} \vdash (\phi \ \& \ X) \rightarrow Z\}$. Clearly, $\phi \in y_0$, but $\psi \notin y_0 \text{ *}_{\text{can}} x$. (Otherwise, $t_{\text{can}} \vdash (\phi \ \& \ X) \rightarrow \psi$ and thus $t_{\text{can}} \vdash X \rightarrow (\phi \rightarrow \psi)$ for some $X \in x$; therefore, $\phi \rightarrow \psi \in y_0 \text{ *}_{\text{can}} x$, a contradiction.) Then, by the Linear Extension Property, we can obtain a linear theory y such that $y_0 \subseteq y$ and $y \text{ *}_{\text{can}} x = \{Z : \text{there is } X \in x \text{ and } t_{\text{can}} \vdash (\phi \ \& \ X) \rightarrow Z\}$; therefore, $\phi \in y$ but $\psi \notin y \text{ *}_{\text{can}} x$.

For (\dagger), we must show

$$x \Vdash_{\text{can}} \phi \ \dagger \ \psi \text{ iff for all } y \in X, \text{ if } y \Vdash_{\text{can}} \phi, \text{ then } x \text{ *}_{\text{can}} y \Vdash_{\text{can}} \psi.$$

Its proof is analogous to that for (\rightarrow). \square

Let us call a model $\mathbf{M}, = (\mathbf{X}, \Vdash_{\text{can}})$ (i.e., $(\mathbf{X}_{\text{can}}, \top_{\text{can}}, \perp_{\text{can}}, t_{\text{can}}, f_{\text{can}}, \leq_{\text{can}}, \text{ *}_{\text{can}}, \Vdash_{\text{can}})$), for **CnHpsUL***, a **CnHpsUL*** model. Then, by Lemma 3.8, the canonically defined $(\mathbf{X}, \Vdash_{\text{can}})$ is a **CnHpsUL*** model. Thus, since, by construction, t_{can} excludes our chosen nontheorem ϕ , and the canonical definition of models agrees with membership, we can state that, for each nontheorem ϕ of **CnHpsUL***, there is a **CnHpsUL*** model in which ϕ is not t_{can} models ϕ . It gives us the weak completeness of **CnHpsUL*** as follows.

Theorem 3.9 (Weak completeness) If $\models_{\text{CnHpsUL*}} \phi$, then \vdash

$\mathbf{CnHpsUL}^* \Phi$.

Furthermore, using Lemma 3.8 and the Linear Extension Property, we can show the strong completeness of $\mathbf{CnHpsUL}^*$ as follows.

Theorem 3.10 (Strong completeness) $\mathbf{CnHpsUL}^*$ is strongly complete w.r.t. the class of all L-frames.

4. Concluding remark

We investigated set-theoretical Kripke-style semantics for weakening-free non-commutative substructural fuzzy logics. As an example, we introduced a set-theoretical Kripke-style semantics for $\mathbf{CnHpsUL}^*$.

References

- Cintula, P. (2006), “Weakly Implicative (Fuzzy) Logics I: Basic properties”, *Archive for Mathematical Logic* 45: pp. 673-704.
- Cintula, P., Horčík, R., and Noguera, C. (2013), “Non-associative substructural logics and their semilinear extensions: axiomatization and completeness properties”, *Review of Symbolic Logic* 6: pp. 394-423.
- Cintula, P., Horčík, R., and Noguera, C. (2015), “The quest for the basic fuzzy logic”, in F. Montagna (ed.) *Petr Hájek on Mathematical Fuzzy Logic*, Dordrecht: Springer, pp. 245-290.
- Cintula, P. and Noguera, C. (2011), A general framework for mathematical fuzzy logic, in P. Cintula, P. Hájek, and C. Noguera (eds.) *Handbook of Mathematical Fuzzy Logic*, vol 1, London: College publications, pp. 103-207.
- Diaconescu, D. (2010), “Kripke-style semantics for non-commutative monoidal t-norm logic”, *Journal of Multiple-Valued Logic and Soft Computing* 16: pp. 247-263.
- Diaconescu, D. and Georgescu, G. (2007), “On the forcing semantics for monoidal t-norm based logic”, *Journal of Universal Computer Science* 13: pp. 1550-1572.
- Esteva, F. and Godo, L. (2001), “Monoidal t-norm based logic: towards a logic for left-continuous t-norms”, *Fuzzy Sets and Systems* 124: pp. 271-288.
- Hájek, P. (2003a), “Fuzzy logics with noncommutative conjunction”, *Journal of Logic and Computation* 13: pp.

469-479.

- Hájek, P. (2003b), “Observations on non-commutative fuzzy logic”, *Soft Computing* 8: pp. 38-43.
- Metcalf, G. and Montagna, F. (2007), “Substructural Fuzzy Logics”, *Journal of Symbolic Logic* 72: pp. 834-864.
- Metcalf, G., Olivetti, N., and Gabbay, D. (2009), *Proof Theory for Fuzzy Logics*, Springer.
- Montagna, F. and Ono, H. (2002), “Kripke semantics, undecidability and standard completeness for Esteva and Godo's Logic MTL_{\forall} ”, *Studia Logica* 71: pp. 227-245.
- Montagna, F. and Sacchetti, L. (2003), “Kripke-style semantics for many-valued logics”, *Mathematical Logic Quarterly* 49: pp. 629-641.
- Montagna, F. and Sacchetti, L. (2004), “Corrigendum to “Kripke-style semantics for many-valued logics”, *Mathematical Logic Quarterly* 50: pp. 104-107.
- Tsinakis, C. and Blount, K. (2003), “The structure of residuated lattices”, *International Journal of Algebra and Computation* 13: pp. 437-461.
- Wang, S. (2013) “Logics for residuated pseudo-uniforms and their residua”, *Fuzzy Sets and Systems* 218: pp. 24-31.
- Wang, S. and Zhao, B. (2009), “HpsUL is not the logic of pseudo-uniforms and their residua”, *Logic Journal of the IGPL* 17: pp. 413-419.
- Yang, E. (2012), “Kripke-style semantics for UL”, *Korean Journal of Logic* 15 (1): pp. 1-15.
- Yang, E. (2014a), “Algebraic Kripke-style semantics for

weakening-free fuzzy logics”, *Korean Journal of Logic* 17: pp. 181-195.

Yang, E. (2014b), “Algebraic Kripke-style semantics for three-valued paraconsistent logic”, *Korean Journal of Logic* 17: pp. 441-460.

Yang, E. (2015a), “Set-theoretic Kripke-style semantics for three-valued paraconsistent logic”, *Korean Journal of Logic* 18: pp. 65-82.

Yang, E. (2015b), “Two kinds of (binary) Kripke-style semantics for three-valued logic”, *Logique et Analyse* 231: pp. 379-396.

Yang, E. (2016), “Algebraic Kripke-style semantics for an extension of HpsUL, CnHpsUL*”, *Korean Journal of Logic* 19: pp. 107-126.

전북대학교 철학과, 비판적사고와논술연구소

Department of Philosophy & Institute of Critical Thinking and Writing, Chonbuk National University

eunsyang@jbnu.ac.kr

CnHpsUL*을 위한 집합 이론적 크립키형 의미론

양 은 석

이 글에서 우리는 약화 없는 비교환적인 퍼지 논리의 비대수적 크립키형 의미론 즉 집합 이론적 크립키형 의미론을 다룬다. 이를 위하여 먼저 우리는 가-유니폼에 기반한 퍼지 논리 HpsUL의 한 확장 체계인 CnHpsUL*을 소개한다. 다음으로 CnHpsUL*을 위한 집합 이론적 크립키형 의미론을 소개한다.

주요어: (집합 이론적) 크립키형 의미론, 대수적 의미론, 퍼지 논리, HpsUL, CnHpsUL*.